



Orange, CA

P: (714) 941-6177 F: (714) 941-6178

Las Vegas, NV

P: (702) 790-4404 F: (702) 790-4406

Allen, TX

P: (469) 257-4200 F: (469) 795-9204

Memphis, TN

P: (901) 560-3580 F: (901) 560-3581

Las Vegas, NV

P: (702) 825-4900

F: (702) 977-8150

Carson City, NV

P: (702) 825-4900

F: (702) 977-8150

Burbank, CA

P: (818) 848-8112

F: (818) 848-8142

Las Vegas, NV

P: (702) 478-5133

F: (702) 478-5401

Neurology Referral Form

Patient Information

Patient Name: _____ DOB: __/__/____
 Sex: Femal Male SS#: ____-____-____ Language: _____
 Allergies: _____ Wt: ____ Kg Lb Ht: ____ Cm In
 Address: _____ Apt/Suite: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Contact: _____ Relation: _____
 Email Address: _____

Prescriber Information

Prescriber Name: _____ NPI: _____
 If Shipping to prescriber: Initial Fill Always Never
 Address: _____ Apt/Suite: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Contact: _____
 Fax: _____ Alt. Fax: _____
 Email Address: _____

INSURANCE INFORMATION (or attach copy of cards)

Medical Plan: _____ Policy #: _____ Policy Holder: _____ Relationship: _____
 Prescription Plan: _____ Policy #: _____ Policy Holder: _____ Relationship: _____

Clinical Information

Access Type: Peripheral PICC PORT
 Has patient previously been on therapy before? No Yes/Last Infusion: _____
 Previously Tried Treatments/Medications for The Condition: _____

Ocrevus (ocrelizumab) 300mg/10mL Vial

Loading Doses
 Infusion 1: 300mg intravenous in 250mL of 0.9% NS.
 Infusion 2: (2 weeks later): 300mg intravenous in 250mL of 0.9% NS.
 Start infusion at 30mL per hour. Increase by 30mL per hour every 30 minutes. Maximum rate: 180mL per hour.
 Maintenance Dose
 Infuse 600mg intravenous in 500mL of 0.9% NS every 6 months (from date of first loading dose).
 Start infusion at 40mL per hour. Increase by 40mL per hour every 30 minutes. Maximum rate: 200mL per hour.
 Refills: _____

Soliris (eculizumab) 300mg/30mL Vial

Loading Doses
 Infuse _____ mg IV every _____ weeks for _____ weeks.
 Maintenance Dose
 Infuse _____ mg IV every _____ weeks. Refills: _____

Ultomiris (ravulizumab) 100mg/mL 1,100mg/11mL Vial OR 300mg/3mL Vial

Loading Doses
 Infuse _____ mg IV on Day 1.
 Maintenance Dose (2 Weeks Later)
 Infuse _____ mg IV every _____ weeks. Refills: _____

Vyvgart (efgartigimod alfa-fcab) 400mg/20mL Vial injection

Loading Doses
 OR Infuse _____ mg/kg IV, 1 time weekly for 4 weeks. Over
 Infuse _____ mg IV, 1 time weekly for 4 weeks. 1 Hour
 Maintenance Dose
 OR Infuse _____ mg/kg IV every _____ weeks. Refills: _____
 Infuse _____ mg IV every _____ weeks.

Tysabri (natalizumab) 300mg/15 mL Vial

- OR Infuse 300mg IV every 4 weeks. Refills: _____
 Infuse 300mg IV every _____ weeks.

Hydration and Pre-Medications

- Hydration: NaCl 0.9% 250mL to infuse at a rate of 250mL/hr IV before & after infusion as hydration is needed.
- Benadryl 25mg capsule: Take 1-2 caps by mouth 30-60 minutes prior to infusion & every 6 hours as needed post infusion.
- Benadryl 50mg ampule: Infuse _____mg slow IVP 30 minutes prior to infusion
- Tylenol 325mg: Take 1-2 tabs by mouth 30-60 minutes prior to infusion & every 6 hours as needed post infusion.
Not to exceed total daily dose of 3000mg.
- Zofran 4mg ODT: Dissolve 1 tab on the tongue every 8 hours as needed for pre & post infusion for nausea and vomiting.
- Solu-Cortef: Infuse _____mg slow IVP 30 minutes prior to infusion
- Solu-Medrol: Infuse _____mg slow IVP 30 minutes prior to infusion

Flushing Protocol

- Sodium Chloride 0.9% 5-10 ml pre and post medications
- Heparin _____ Units/mL OR _____ mL as needed

Anaphylaxis Protocol

MILD infusion reactions: (Slow infusion rate by 50% until symptoms resolve. Resume at previous rate as tolerated)
Diphenhydramine 25mg caps #4 Sig: Take 1-2 capsules PO Q6H PRN for infusion reactions, NTE total daily dose of 400mg (16 caps/day).

MODERATE infusion reactions: (Stop Infusion, resume at 50% rate when symptoms resolve)
Diphenhydramine 50mg/ml vial #1 Sig: Inject 1ml (=50mg) IV by slow IV push over 3-5 mins PRN for anaphylaxis reaction.
May repeat every 4 hrs, NTE 400mg/24hrs

SEVERE ANAPHYLAXIS *CALL 911* (Stop infusion and remove tubing from access device to prevent further administration)

1. Epinephrine 0.3mg Auto Injector #2 Sig: Remove cap then inject auto-injector intramuscularly in the thigh area and hold for 2 to 10 seconds as needed for anaphylaxis reaction, may administer through clothing if necessary. May repeat in 5-15-minutes.
2. Initiate 0.9% NaCl 500 mL IV x 1
3. Administer CPR if needed until EMS arrives

Ancillary Supplies : As needed for proper administration and disposal of medication

Skilled Nursing Visits: As needed for IV access, administration and proper clinical monitoring

Administration procedures to be followed per pharmacy protocol.

Prescription will be filled with generic (if available) unless prescriber writes "DAW" (dispense as written): _____

Deliver to: Home Office Infusion Suite Other: _____

If shipped to prescriber's office or infusion clinic, prescriber accepts on behalf of patient for administration in office or infusion clinic.
By signing this order, I, the prescriber, certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber's Signature _____

Date: __/__/__

* Prescriber Authorization: I authorize this pharmacy and its representatives to act as my authorized agent to secure coverage and initiate the insurance prior authorization process for my patient(s), and to sign any necessary forms on my behalf as my authorized agent, including the receipt of any required prior authorization forms and the receipt and submission of patient lab values and other patient data. In the event that this pharmacy determines that it is unable to fulfill this prescription, I further authorize this pharmacy to forward this information and any related materials related to coverage of the product to another pharmacy of the patient's choice or in the patient's insurer's provider network.